

# percept™

cell-free DNA prenatal test

## What is the percept™ cfDNA prenatal test?


*percept*™ is an Australian non-invasive prenatal test (NIPT) offered by Victorian Clinical Genetics Services (VCGS). This test uses cell-free fetal DNA (cfDNA) found in maternal blood to identify pregnancies with an increased chance of chromosome conditions. It is the most accurate type of prenatal screening test for Down syndrome.

*percept*™ by VCGS is Australasia's most advanced non-invasive prenatal test: *simple, fast, reliable*

## When can I have this test?


The *percept* blood test can be performed from 10 weeks of pregnancy onwards, as confirmed by an acceptable dating method like ultrasound. Test results are usually available within 3-5 working days from the time we receive the sample.

1. You can choose *percept* as your initial screening test from 10 weeks. VCGS recommends that you also consider having a 12-13 week ultrasound to check the growth of your baby.



2. Alternatively, you may choose to have *percept* after combined first trimester screening (CFTS). CFTS includes an ultrasound and a blood test.

If you are not reassured by the results of your CFTS, further screening with *percept* may be helpful.



## What will this test tell me?

Conditions screened	Detection Rate
<b>Trisomies</b>	
Down syndrome (trisomy 21)	>99%
Edwards syndrome (trisomy 18)	>98%
Patau syndrome (trisomy 13)	>98%
Rare trisomies	LD*
<b>Sex chromosome conditions</b>	
Turner syndrome (monosomy X)	>95%
Klinefelter syndrome (XXY)	LD*
Triple X (XXX)	LD*
XYY	LD*
<b>Fetal sex</b>	
Male XY, Female XX	>99%


\*Limited data available

- ✓ Results within 3-5 working days (from receipt)
- ✓ Available from 10 weeks of pregnancy
- ✓ IVF pregnancy
- ✓ Egg donor pregnancy
- ✓ Twin pregnancy (trisomies only)

You should discuss your options with your healthcare provider. Genetic counselling is available through VCGS. Please see [vcgs.org.au/perceptNIPT](http://vcgs.org.au/perceptNIPT) for more information.

## How does *percept*<sup>TM</sup> work?

During pregnancy, some DNA from the placenta crosses into your bloodstream. *percept* tests this cell-free DNA in your blood to look for certain chromosome conditions that may be affecting your pregnancy.



## When should I consider diagnostic testing instead of NIPT?

NIPT identifies a limited range of chromosome conditions. In some situations, your doctor may recommend a diagnostic test such as chorionic villus sampling (CVS) or amniocentesis with chromosome analysis. This provides more detailed and accurate information about your pregnancy. These situations may include:

- If the ultrasound shows a fetal abnormality.
- If the combined first trimester screening risk is very high (>1 in 10).
- If the nuchal translucency is increased at the 12 week ultrasound.

**NIPT is an advanced screening test but is NOT a replacement for diagnostic testing through CVS or amniocentesis.**

## How do I arrange this test?



Discuss *percept* with your healthcare provider.



A test request form will be completed by your healthcare provider.



Have your blood collected. For collection sites, see the Mater Pathology website.



Most results available within 3-5 working days (from receipt of sample).

## How are *percept*<sup>TM</sup> results reported?

### Low risk

Most results are reported as low risk. This means it is very unlikely (less than 1 in 10,000 chance) that your pregnancy is affected by the specific conditions screened. However, other conditions or abnormalities may still be present. We recommend all patients have an 18-20 week ultrasound.

### High risk

A high risk result means that there is an increased chance of the chromosome condition in your pregnancy. **CVS or amniocentesis with chromosome analysis is strongly recommended to confirm a high risk result.** In some cases the diagnostic test will show your baby is not affected by the condition. In this case, the NIPT result is considered a 'false-positive' result.

### No result

In some cases, no result is obtained. This is very uncommon (<1.0% of samples). If this occurs the laboratory will request a repeat specimen for testing at no additional charge.

## What does *percept*<sup>TM</sup> cost?

This test is not covered by private health insurance or Medicare (for Australian patients). Please see [pathology.mater.org.au](http://pathology.mater.org.au) for current pricing.

## Contact details

Mater Pathology  
Level 6, Mater Hospital Brisbane  
Raymond Terrace  
South Brisbane, QLD 4101

P (07) 3163 8500

W [pathology.mater.org.au](http://pathology.mater.org.au)

F (07) 3163 8338

E [pathology.enquiries@mater.org.au](mailto:pathology.enquiries@mater.org.au)

## What is Down syndrome (trisomy 21)?

Down syndrome is the most common chromosomal condition seen in children and adults. It is caused by an extra copy of chromosome 21. People with Down syndrome have some degree of intellectual disability and may have other health and developmental challenges.

## What is Edwards syndrome (trisomy 18)?

Edwards syndrome is less common than Down syndrome and is caused by an extra chromosome 18. Pregnancies with trisomy 18 usually miscarry and babies that are liveborn rarely survive for long and will have significant physical and intellectual problems.

## What is Patau syndrome (trisomy 13)?

Patau syndrome is even less common than Down syndrome and Edwards syndrome. It is caused by an extra copy of chromosome 13. Pregnancies with trisomy 13 usually miscarry and babies that are liveborn rarely survive for long.

## What are sex chromosome conditions?

The X and Y chromosomes are called the sex chromosomes because they determine if a baby will be male or female. Sometimes there are extra or missing copies of these chromosomes. Development may be affected in a variety of ways, depending on the chromosome condition.

## What are rare trisomies?

Rare trisomies are caused by three copies of any chromosome other than 21, 18, 13, X or Y. These are rarely seen at birth and can be associated with an increased chance of an early miscarriage, growth problems in the fetus and premature delivery. Babies that are liveborn may have a chromosomal condition.

**NOTE:** *percept* may be used to screen a pregnancy where one parent is a known translocation carrier. Testing only performed with prior arrangement.

